

On a generic defining function of projective motion in the rigid 6-dimensional h -spaces ¹

Zolfira Zakirova²

Abstract

In this note we find a generic defining function of projective motion in the 6-dimensional rigid h -space.

1 Introduction

In the series of works [Z]–[Z6], we investigate the 6-dimensional pseudo-Riemannian space $V^6(g_{ij})$ with signature $[+ + - - - -]$, which admit projective motions, i.e. continuous transformation groups preserving geodesics. The general method of determining pseudo-Riemannian manifolds that admit some nonhomothetic projective group G_r has been developed by A.V.Aminova [A95]. A.V.Aminova has classified all the Lorentzian manifolds of dimension ≥ 3 that admit nonhomothetic projective or affine infinitesimal transformations. In each case, there were determined the corresponding maximal projective and affine Lie algebras. This problem is not solved for pseudo-Riemannian spaces with arbitrary signature. In the papers [Z]–[Z5], we found 6-dimensional rigid h -spaces of the [2211], [321], [33], [411] and [51] types:

the metric of the h -space of the [2211] type

$$g_{ij}dx^i dx^j = e_2(f_4 - f_2)^2 \Pi_\sigma(f_\sigma - f_2) \{2A dx^1 dx^2 - A^2 \Sigma_1(dx^2)^2\} + \quad (1)$$

$$e_4(f_2 - f_4)^2 \Pi_\sigma(f_\sigma - f_4) \{2\tilde{A} dx^3 dx^4 - \tilde{A}^2 \Sigma_2(dx^4)^2\} + \sum_{\sigma} e_\sigma \Pi'_i(f_i - f_\sigma)(dx^\sigma)^2,$$

¹The research was partially supported by the RFBR grant 04-02-16538-a.

²Kazan State University, e-mail: Zolfira.Zakirova@soros.ksu.ru

where

$$A = \epsilon x^1 + \theta(x^2), \quad \tilde{A} = \tilde{\epsilon} x^3 + \omega(x^4), \quad (2)$$

$$\Sigma_1 = 2(f_4 - f_2)^{-1} + \sum_{\sigma} (f_{\sigma} - f_2)^{-1}, \quad \Sigma_2 = 2(f_2 - f_4)^{-1} + \sum_{\sigma} (f_{\sigma} - f_4)^{-1},$$

$f_1 = f_2 = \epsilon x^2$, $f_3 = f_4 = \tilde{\epsilon} x^4 + a$, $\epsilon, \tilde{\epsilon} = 0, 1$, $\epsilon \neq \tilde{\epsilon}$, a is a constant nonzero whenever $\tilde{\epsilon} = 0$, $f_{\sigma}(x^{\sigma})$, $\theta(x^2)$, $\omega(x^4)$ are arbitrary functions, $e_i = \pm 1$.

the metric of the h -space of the [321] type

$$g_{ij} dx^i dx^j = e_3 (f_5 - \epsilon x^3)^2 (f_6 - \epsilon x^3) \{ (dx^2)^2 + 4A dx^1 dx^3 + \quad (3)$$

$$2(\epsilon x^1 - 2A \Sigma_1) dx^2 dx^3 + ((\epsilon x^1)^2 - 4\epsilon x^1 A \Sigma_1 + 4A^2 \Sigma_3) (dx^3)^2 \} +$$

$$e_5 (\epsilon x^3 - f_5)^3 (f_6 - f_5) \{ 2\tilde{A} dx^4 dx^5 - \Sigma_4 \tilde{A}^2 (dx^5)^2 \} + e_6 (f_5 - f_6)^2 (f_5 - f_6)^3 (dx^6)^2,$$

where

$$A = \epsilon x^2 + \theta(x^3), \quad \tilde{A} = \tilde{\epsilon} x^4 + \omega(x^5), \quad (4)$$

$$\Sigma_1 = (f_6 - \epsilon x^3)^{-1} + 2(f_5 - \epsilon x^3)^{-1}, \quad \Sigma_2 = (f_6 - \epsilon x^3)^{-2} + 2(f_5 - \epsilon x^3)^{-2},$$

$$2\Sigma_3 = (\Sigma_1)^2 - \Sigma_2, \quad \Sigma_4 = 3(\epsilon x^3 - f_5)^{-1} + (f_6 - f_5)^{-1},$$

$f_1 = f_2 = f_3 = \epsilon x^3$, $f_4 = f_5 = \tilde{\epsilon} x^5 + a$, $\theta(x^3)$, $\omega(x^5)$, $f_6(x^6)$ are arbitrary function, $\epsilon, \tilde{\epsilon} = 0, 1$, $\epsilon \neq \tilde{\epsilon}$, a is a constant nonzero whenever $\tilde{\epsilon} = 0$, $e_3, e_5, e_6 = \pm 1$.

the metric of the h -space of the [33] type

$$g_{ij} dx^i dx^j = e_3 (f_6 - f_3)^3 \{ (dx^2)^2 + 4A dx^1 dx^3 + \quad (5)$$

$$2(\epsilon x^1 - 2A \Sigma_1) dx^2 dx^3 + ((\epsilon x^1)^2 - 4\epsilon x^1 A \Sigma_1 + 4A^2 \Sigma_2) (dx^3)^2 \} +$$

$$e_6 (f_3 - f_6)^3 \{ (dx^5)^2 + 4\tilde{A} dx^4 dx^6 + 2(\tilde{\epsilon} x^4 + 2\tilde{A} \Sigma_1) dx^5 dx^6 + ((\tilde{\epsilon} x^4)^2 + 4\tilde{\epsilon} x^4 \tilde{A} \Sigma_1 + 4\tilde{A}^2 \Sigma_2) (dx^6)^2 \},$$

where

$$A = \epsilon x^2 + \theta(x^3), \quad \tilde{A} = \tilde{\epsilon} x^4 + \omega(x^6), \quad (6)$$

$$\Sigma_1 = 3(f_6 - f_3)^{-1}, \quad \Sigma_2 = 3(f_6 - f_3)^{-2},$$

$f_1 = f_2 = f_3 = \epsilon x^3$, $f_4 = f_5 = f_6 = \tilde{\epsilon} x^6 + a$, $\epsilon, \tilde{\epsilon} = 0, 1$, $\epsilon \neq \tilde{\epsilon}$, a is a constant nonzero whenever $\tilde{\epsilon} = 0$, $e_i = \pm 1$, $\theta(x^3)$, $\omega(x^6)$ are arbitrary functions.

the metric of the h -space of the [411] type

$$g_{ij} dx^i dx^j = e_4 \Pi_{\sigma} (f_{\sigma} - \epsilon x^4) \{ 6A dx^1 dx^4 + 2dx^2 dx^3 + 2(2\epsilon x^2 - 3A \Sigma_1) dx^2 dx^4 - \quad (7)$$

$$\begin{aligned} & \Sigma_1(dx^3)^2 + 2(\epsilon x^1 - 2\epsilon x^2 \Sigma_1)dx^3 dx^4 + 4((\epsilon x^2)^2 \Sigma_1 + \epsilon^2 x^1 x^2 - \frac{3}{2}\epsilon x^1 A \Sigma_1)(dx^4)^2 \} + \\ & 3A dx^3 dx^4 + 12\epsilon x^2 A (dx^4)^2 + \sum_{\sigma} e_{\sigma} \Pi'_i (f_i - f_{\sigma})(dx^{\sigma})^2, \end{aligned}$$

where

$$A = \epsilon x^3 + \theta(x^4), \quad \Sigma_1 = (f_5 - \epsilon x^4)^{-1} + (f_6 - \epsilon x^4)^{-1}, \quad (8)$$

$f_1 = f_2 = f_3 = f_4 = \epsilon x^4$, $f_5(x^5)$, $f_6(x^6)$, $\theta(x^4)$ are arbitrary functions, $e_4, e_5, e_6 = \pm 1$, $\epsilon = 0, 1$

the metric of the h -space of the [51] type

$$\begin{aligned} g_{ij} dx^i dx^j = & e_5 (f_6 - \epsilon x^5) \{ 8A dx^1 dx^5 + 2dx^2 dx^4 + \\ & 2(3\epsilon x^3 - 4A \Sigma_1) dx^2 dx^5 + (dx^3)^2 - 2\Sigma_1 dx^3 dx^4 + 2(2\epsilon x^2 - 3\epsilon x^3 \Sigma_1) dx^3 dx^5 + \\ & 2(\epsilon x^1 - 2\epsilon x^2 \Sigma_1) dx^4 dx^5 + 4(3/2\epsilon x^1 \epsilon x^3 + (\epsilon x^2)^2 - 2\epsilon x^1 A \Sigma_1 - \\ & 3\epsilon x^2 \epsilon x^3 \Sigma_1)(dx^5)^2 \} + e_6 (\epsilon x^5 - f_6)^5 (dx^6)^2, \end{aligned} \quad (9)$$

where

$$A = \epsilon x^4 + \theta(x^5), \quad \Sigma_1 = (f_6 - \epsilon x^5)^{-1}, \quad (10)$$

$\theta(x^5)$, $f_6(x^6)$ are arbitrary functions, $f_1 = f_2 = f_3 = f_4 = f_5 = \epsilon x^5$, $e_5, e_6 = \pm 1$, $\epsilon = 0, 1$.

The defining function of projective motion in the 6-dimensional rigid h -spaces is

$$\varphi = \frac{1}{2} \sum_{i=1}^6 f_i. \quad (11)$$

In the paper [Z6], we determined conditions of the constant curvature of the 6-dimensional rigid h -spaces:

for the h -space of the [2211] type

$$\rho_p - \rho_{\sigma p} = \rho_p - \rho_{pq} = \epsilon = \tilde{\epsilon} = 0 \quad (p \neq q, p, q = 2, 4, \sigma = 5, 6), \quad (12)$$

for the h -space of the [321] type

$$f'_6 = \epsilon = \tilde{\epsilon} = 0, \quad (13)$$

for the h -space of the [33] type

$$\epsilon = \tilde{\epsilon} = 0, \quad (14)$$

for the h -space of the [411] type

$$\rho_p - \rho_{\sigma p} = \epsilon = \gamma_1 = \gamma_2 = 0 \quad (\sigma = 5, 6, p = 4), \quad (15)$$

for the h -space of the [51] type

$$f'_6 = \epsilon = 0, \quad (16)$$

where ρ_p , $\rho_{\sigma p}$, ρ_{pq} , γ_1 and γ_2 are determined by formulas

$$\rho_p = -\frac{1}{4} \sum_{\sigma} \frac{(f'_{\sigma})^2}{(f_{\sigma} - f_p)^2 g_{\sigma\sigma}}, \quad \rho_{pq} = -\frac{1}{4} \sum_{\sigma} \frac{(f'_{\sigma})^2}{(f_{\sigma} - f_p)(f_{\sigma} - f_q) g_{\sigma\sigma}}, \quad (17)$$

$$\begin{aligned} \rho_{\sigma p} = & -\frac{1}{4} \frac{(f'_{\sigma})^2}{(f_{\sigma} - f_p) g_{\sigma\sigma}} \left\{ \frac{2f''_{\sigma}}{(f'_{\sigma})^2} - \frac{1}{f_{\sigma} - f_p} + \sum_{i, i \neq \sigma} (f_i - f_{\sigma})^{-1} \right\} \\ & - \frac{1}{4} \sum_{\gamma, \gamma \neq \sigma} \frac{(f'_{\gamma})^2}{(f_{\gamma} - f_p)(f_{\gamma} - f_{\sigma}) g_{\gamma\gamma}}. \\ \gamma_1 = & -\frac{1}{4} \sum_{\sigma} \frac{(f'_{\sigma})^2}{(f_{\sigma} - f_4)^3 g_{\sigma\sigma}}, \quad \gamma_2 = -\frac{1}{4} \sum_{\sigma} \frac{(f'_{\sigma})^2}{(f_{\sigma} - f_4)^4 g_{\sigma\sigma}}. \end{aligned} \quad (18)$$

The results obtained in the papers [Z]-[Z6] are necessary for investigating properties of the defining function of projective motion in the rigid 6-dimensional h -spaces.

2 Generic defining function of projective motion in the rigid 6-dimensional h -spaces

In this section we prove the following theorem that manifestly gives any defining function of projective motion.

Theorem. *Any defining function of projective motion in the rigid h -spaces of nonconstant curvature can be presented as $\phi = a_1 \varphi$, where a_1 is a constant and φ is determined by (11).*

Proof. Suppose we are given with a vector field ξ^i that gives the projective motion with the defining function ϕ in the rigid h -spaces. Then, for the tensor $b_{ij} = L_{\xi} g_{ij}$ the following Eisenhart equations [E] are fulfilled

$$b_{ij,k} = 2g_{ij}\phi_{,k} + g_{ik}\phi_{,j} + g_{jk}\phi_{,i}. \quad (19)$$

Their integrability conditions are

$$b_{mi}R_{jkl}^m + b_{mj}R_{ikl}^m = g_{ik}\phi_{,jl} + g_{jk}\phi_{,il} - g_{li}\phi_{,jk} - g_{lj}\phi_{,ik}. \quad (20)$$

Let us show that for any projective motion in the rigid h -spaces the following conditions are satisfied

$$b_{\alpha\beta} = 0, \quad \phi_{,\alpha\beta} = 0, \quad (21)$$

α, β being indices of vanishing components of the metric $g_{\alpha\beta}$ of the h -spaces of the [2211], [321], [33], [411] and [51] type.

For the h -space of the [2211] type, the set $(ijkl) = (3112), (3314), (3\sigma 1\sigma), (1334), (1132), (1\sigma 3\sigma)$. From (20), one finds

$$b_{13}R_{112}^1 = -g_{12}\phi_{,13}, \quad b_{13}R_{314}^1 = -g_{34}\phi_{,13}, \quad b_{13}R_{\sigma 1\sigma}^1 = -g_{\sigma\sigma}\phi_{,13}, \quad (22)$$

$$b_{13}R_{334}^3 = -g_{34}\phi_{,13}, \quad b_{13}R_{132}^3 = -g_{12}\phi_{,13}, \quad b_{13}R_{\sigma 3\sigma}^3 = -g_{\sigma\sigma}\phi_{,13}, \quad (23)$$

From (22) it follows that

$$b_{13}(\chi_2 - \rho_{24}) = b_{13}(\chi_2 - \rho_{\sigma 2}) = 0,$$

and from (23)

$$b_{13}(\chi_4 - \rho_{24}) = b_{13}(\chi_4 - \rho_{\sigma 4}) = 0.$$

If $b_{13} \neq 0$, then $\chi_p - \rho_{\sigma p} = 0$ ($p = 2, 4$). Since we consider h -spaces of the nonconstant curvature, one necessarily puts $b_{13} = 0$, hence $\phi_{,13} = 0$. Similarly one can get other equalities (21).

In the case of the h -space of the [321] type, when $(ijkl) = (1212), (1415), (1515), (1616), (2212), (1123), (2415), (4125)$, equations (20) leads to the following identities

$$b_{11}R_{212}^1 = -g_{22}\phi_{,11}, \quad b_{11}R_{123}^1 = b_{11}R_{425}^1 = 0, \quad (24)$$

$$b_{11} \frac{R_{415}^1}{g_{45}} = b_{11} \frac{R_{515}^1}{g_{55}} = b_{11} \frac{R_{616}^1}{g_{66}} = \phi_{,11}, \quad (25)$$

hence

$$b_{11}(\chi_3 - \rho_{63}) = b_{11}(\chi_3 - \rho_{53}) = b_{11}(\chi_3 - \rho_{53} - \frac{\tilde{\epsilon}\omega'}{(f_5 - f_3)\tilde{A}g_{55}}) = 0.$$

If $b_{11} \neq 0$, then $\chi_3 - \rho_{63} = \chi_3 - \rho_{53} = \tilde{\epsilon} = 0$. From this it follows that $\epsilon = \rho_3 - \rho_{53} = 0$, and $f_6' = 0$. Therefore, $\chi_5 - \rho_{53} = \chi_5 - \rho_{65} = \gamma = 0$, this leads to the constant curvature h -space. In such a manner, $b_{11} = \phi_{,11} = 0$. Similarly one can get other equalities (21).

To prove that $b_{11} = \phi_{,11} = 0$ for the h -space of the [33] type, set $(ijkl) = (1212), (1416)$ and (1516) in (20). Then, one obtains

$$b_{11}R_{212}^1 = -g_{22}\phi_{,11}, \quad b_{11}R_{416}^1 = -g_{46}\phi_{,11}, \quad (26)$$

$$b_{11}(\frac{R_{212}^1}{g_{22}} - \frac{R_{516}^1}{g_{56}}) = 0. \quad (27)$$

Since $R_{416}^1 = 0$ and $g_{46} \neq 0$, we get $\phi_{,11} = 0$. Then, from (27) it follows that $b_{11} = 0$, since $R_{212}^1 \neq 0$. Similarly one can get other equalities (21).

Putting $(ijkl) = (1213), (1313), (1\sigma 1\sigma), (3312), (1314)$ in the case of the h -space of the [411] type, one gets

$$b_{11}R_{213}^1 = -g_{23}\phi_{,11}, \quad b_{11}R_{313}^1 = -g_{33}\phi_{,11}, \quad (28)$$

$$b_{11}R_{\sigma 1\sigma}^1 = -g_{\sigma\sigma}\phi_{,11}, \quad (29)$$

$$b_{13}R_{312}^1 = -g_{23}\phi_{,13}, \quad b_{11}R_{314}^1 + b_{13}R_{114}^1 = -g_{14}\phi_{,13} - g_{34}\phi_{,11}. \quad (30)$$

Since

$$\begin{aligned} R_{213}^1 &= R_{312}^1 = g_{23}\rho_4, & R_{114}^1 &= g_{14}\rho_4, \\ R_{313}^1 &= g_{33}\rho_4 + \gamma_1 g_{23} + \frac{2\epsilon^2}{9A^2}, & R_{\sigma 1\sigma}^1 &= g_{\sigma\sigma}\rho_{\sigma 4}, \\ R_{314}^1 &= g_{34}\rho_4 + \gamma_1 g_{24} + \gamma_2 g_{14} + \frac{2\epsilon}{3A^2}(\theta' - \frac{4\epsilon^2 x^2}{3}), \end{aligned}$$

then, from (28) and (29), we have

$$b_{11}(\rho_4 - \rho_{\sigma 4}) = b_{11}(\rho_4 - \rho_{\sigma 4} + \gamma_3 \frac{g_{23}}{g_{33}} + \frac{2\epsilon^2}{9A^2 g_{23}}) = 0.$$

From (29) and (30), one finds

$$b_{11}(\rho_4 - \rho_{\sigma 4} + \gamma_3 \frac{g_{24}}{g_{34}} + \gamma_4 \frac{g_{14}}{g_{34}} + \frac{2\epsilon}{3A^2 g_{34}}(\theta' - \frac{4\epsilon^2 x^2}{3})) = 0.$$

Let us $b_{11} \neq 0$. From the last relations it follows that $\rho_4 - \rho_{\sigma 4} = \epsilon = \gamma_1 = \gamma_2 = 0$, which are necessary and sufficient conditions of the constant curvature of these h -spaces, therefore, $b_{11} = 0$, and, hence, $\phi_{,11} = 0$. Similarly one can get other equalities (21).

For the h -space of the [51] type from integrability conditions one can similarly get equalities (21).

Consider equation (19). Putting $(ijk) = (1\sigma\sigma), (3\sigma\sigma), (\sigma\tau\sigma), (\tau\sigma\tau)$ ($\sigma, \tau = 5, 6, \sigma \neq \tau$) for the h -space of the [2211] type, we determine

$$\phi_{,1} = 0, \quad \phi_{,3} = 0, \quad (31)$$

$$\phi_{,\sigma} = f'_\sigma P_{\sigma\tau}, \quad \phi_{,\tau} = f'_\tau P_{\sigma\tau}, \quad P_{\sigma\tau} \equiv \frac{1}{2} \frac{b_{\tau\tau} g_{\tau\tau}^{-1} - b_{\sigma\sigma} g_{\sigma\sigma}^{-1}}{f_\tau - f_\sigma}, \quad (\sigma \neq \tau). \quad (32)$$

From this it follows that $f'_\sigma \phi_{,\tau} = f'_\tau \phi_{,\sigma}$. Therefore, from the condition $\phi_{,\sigma\tau} = 0$, one gets

$$\partial_{\sigma\tau}\phi = 0. \quad (33)$$

Using relations obtained from (19) with $(ijk) = (1\tau 2), (2\tau\tau), (\tau\tau\tau)$, we find

$$\phi_{,\tau} = \frac{1}{2} \frac{f'_\tau}{f_\tau - f_2} \left(\frac{b_{\tau\tau}}{g_{\tau\tau}} - \frac{b_{12}}{g_{12}} \right), \quad (34)$$

$$\phi_{,2} = \frac{\epsilon}{f_\tau - f_2} \left(\frac{b_{\tau\tau}}{g_{\tau\tau}} - \frac{b_{12}}{g_{12}} \right), \quad (35)$$

$$\frac{\partial_\tau b_{\tau\tau}}{g_{\tau\tau}} = -f'_\tau \sum_{i,j \neq \tau} (f_i - f_\tau)^{-1} \frac{b_{\tau\tau}}{g_{\tau\tau}} + 4\phi_{,\tau}. \quad (36)$$

From (34) and (35), we have

$$2\epsilon\phi_{,\tau} = f'_\tau \phi_{,2}. \quad (37)$$

These relations and conditions $\phi_{,2\tau} = 0$ give

$$\partial_{2\tau}\phi = 0. \quad (38)$$

Differentiating (34) w.r.t. x^τ and taking into account (36), one obtains

$$f'_\tau \partial_\tau \phi_{,\tau} = f''_\tau \phi_{,\tau}. \quad (39)$$

Now, from (19) with $(ijk) = (142), (234)$, one finds

$$\phi_{,2} = \frac{\tilde{\epsilon}}{f_4 - f_2} \left(\frac{b_{34}}{g_{34}} - \frac{b_{12}}{g_{12}} \right), \quad \phi_{,4} = \frac{\epsilon}{f_4 - f_2} \left(\frac{b_{34}}{g_{34}} - \frac{b_{12}}{g_{12}} \right), \quad (40)$$

hence, $\epsilon\phi_{,4} = \tilde{\epsilon}\phi_{,2}$ and, since $\phi_{,24} = 0$, one obtains $\partial_{24}\phi = 0$.

Assuming $(ijk) = (3\tau 4), (4\tau\tau)$ in equation (19), one obtains

$$2\tilde{\epsilon}\phi_{,\tau} = f'_\tau \phi_{,4}, \quad (41)$$

therefore, taking into account $\phi_{,4\tau} = 0$, we have $\partial_{4\tau}\phi = 0$. If not all $f'_\tau = 0$, then, from (37), (39) and (41), one finds

$$\phi_{,\tau} = \frac{1}{2} a_1 f'_\tau, \quad \phi_{,2} = a_1 \epsilon, \quad \phi_{,4} = a_1 \tilde{\epsilon}. \quad (42)$$

The first equation of (42) is correct whenever $f'_\tau = 0$ due to relation (34). Integrating equation (42), one obtains

$$\phi = \frac{1}{2} a_1 \sum_{i=1}^6 f_i = a_1 \varphi, \quad f_1 = f_2 = \epsilon x^2, \quad f_3 = f_4 = \tilde{\epsilon} x^4 + a, \quad (43)$$

where φ determined by formula (11). If all $f'_\tau = 0$, then, from (20) with $(ijkl) = (\tau 2 \tau 2), (\tau 4 \tau 4)$, one finds $\partial_{22}\phi = \partial_{44}\phi = 0$, and, therefore, (43) is correct.

For the h -space of the [321] type, putting in equation (19) $(ijk) = (166), (266), (466)$, one obtains $\phi_{,1} = \phi_{,2} = \phi_{,4} = 0$.

Equation (19) with $(ijk) = (163), (366), (666)$ leads to

$$\phi_{,6} = \frac{1}{2} \frac{f'_6}{f_6 - f_3} \left(\frac{b_{66}}{g_{66}} - \frac{b_{13}}{g_{13}} \right), \quad (44)$$

$$\phi_{,3} = \frac{3}{2} \frac{\epsilon}{f_6 - f_3} \left(\frac{b_{66}}{g_{66}} - \frac{b_{13}}{g_{13}} \right), \quad (45)$$

$$\frac{\partial_6 b_{66}}{g_{66}} = -f'_6 \sum_{i,j \neq 6} (f_i - f_6)^{-1} \frac{b_{66}}{g_{66}} + 4\phi_{,6}, \quad (46)$$

hence

$$f'_6 \partial_6 \phi_{,6} = f''_6 \phi_{,6}, \quad (47)$$

$$3\epsilon \phi_{,6} = f'_6 \phi_{,3}. \quad (48)$$

Since $\phi_{,36} = 0$, then, from equation (48), it follows that $\partial_{36}\phi = 0$. In case of $(ijk) = (465), (566)$, from (19) we have

$$\phi_{,6} = \frac{1}{2} \frac{f'_6}{f_6 - f_2} \left(\frac{b_{66}}{g_{66}} - \frac{b_{12}}{g_{12}} \right), \quad (49)$$

$$\phi_{,2} = \frac{\epsilon}{f_6 - f_2} \left(\frac{b_{66}}{g_{66}} - \frac{b_{12}}{g_{12}} \right), \quad (50)$$

and, therefore,

$$2\epsilon \phi_{,6} = f'_6 \phi_{,2}. \quad (51)$$

Substituting $(ijk) = (345), (513)$ in (19), one finds

$$\phi_{,3} = \frac{3}{2} \frac{\epsilon}{f_5 - f_3} \left(\frac{b_{45}}{g_{45}} - \frac{b_{13}}{g_{13}} \right), \quad \phi_{,5} = \frac{\tilde{\epsilon}}{f_5 - f_3} \left(\frac{b_{45}}{g_{45}} - \frac{b_{13}}{g_{13}} \right), \quad (52)$$

therefore,

$$3\epsilon \phi_{,5} = 2\tilde{\epsilon} \phi_{,3}. \quad (53)$$

From this relation, taking into account $\phi_{,35} = 0$, we find $\partial_{35}\phi = 0$. If $f'_6 \neq 0$, then, integrating the equations (47), (48) and (53), one obtains

$$\phi = \frac{1}{2} a_1 \sum_{i=1}^6 f_i = a_1 \varphi, \quad f_1 = f_2 = f_3 = \epsilon x^3, \quad f_4 = f_5 = \tilde{\epsilon} x^5 + a, \quad (54)$$

where φ is determined by formula (11).

On the contrary, from equation (20) with $(ijkl) = (3312), (3636), (6565)$ one can obtain $\partial_{33}\phi = \partial_{55}\phi = 0$ which leads to the desired result.

From (19) with $(ijk) = (166), (266), (433), (533), (346), (613)$ in the case of the h -space of the [33] type, one obtains the following relations

$$\begin{aligned} \phi_{,1} &= \phi_{,2} = \phi_{,4} = \phi_{,5} = 0, \\ \phi_{,3} &= \frac{3}{2} \frac{\epsilon}{f_6 - f_3} \left(\frac{b_{46}}{g_{46}} - \frac{b_{13}}{g_{13}} \right), \quad \phi_{,6} = \frac{3}{2} \frac{\tilde{\epsilon}}{f_6 - f_3} \left(\frac{b_{46}}{g_{46}} - \frac{b_{13}}{g_{13}} \right). \end{aligned}$$

Hence

$$\tilde{\epsilon}\phi_{,3} = \epsilon\phi_{,6}. \quad (55)$$

Therefore, from the condition $\phi_{,36} = 0$, one obtains $\partial_{36}\phi = 0$. If $\tilde{\epsilon} \neq 0$, then, differentiating relation (55) w.r.t. x^3 , we have $\partial_{33}\phi = 0$. If $\tilde{\epsilon} = 0$, then $\epsilon \neq 0$, since we consider the space of nonconstant curvature. In case when $\tilde{\epsilon} = 0$ and $\epsilon \neq 0$, the relation $\partial_{33}\phi = 0$ follows from (20) with $(ijkl) = (3312), (3436)$. Similarly one can get that $\partial_{66}\phi = 0$. Integrating the obtained equations, one finds

$$\phi = \frac{1}{2} a_1 \sum_{i=1}^6 f_i = a_1 \varphi, \quad f_1 = f_2 = f_3 = \epsilon x^3, \quad f_3 = f_4 = f_6 = \tilde{\epsilon} x^6 + a, \quad (56)$$

where φ is determined by formula (11).

For the h -space of the [411] type from (19) with $(ijk) = (1\tau\tau), (2\tau\tau), (3\tau\tau)$ ($\tau = 5, 6$), one obtains

$$\phi_{,1} = \phi_{,2} = \phi_{,3} = 0.$$

For this h -space relations (32) are also correct. From these it follows that $\partial_{\sigma\tau}\phi = 0$ ($\sigma, \tau = 5, 6, \sigma \neq \tau$). Using the obtained results, from (19) $(ijk) = (1\tau 4), (4\tau\tau), (\tau\tau\tau)$, we find

$$\phi_{,\tau} = \frac{1}{2} \frac{f'_\tau}{f_\tau - f_4} \left(\frac{b_{\tau\tau}}{g_{\tau\tau}} - \frac{b_{14}}{g_{14}} \right), \quad (57)$$

$$\phi_{,4} = 2 \frac{\epsilon}{f_\tau - f_4} \left(\frac{b_{\tau\tau}}{g_{\tau\tau}} - \frac{b_{14}}{g_{14}} \right), \quad (58)$$

$$\frac{\partial_\tau b_{\tau\tau}}{g_{\tau\tau}} = -f'_\tau \sum_{i,j \neq \tau} (f_i - f_\tau)^{-1} \frac{b_{\tau\tau}}{g_{\tau\tau}} + 4\phi_{,\tau}, \quad (59)$$

hence,

$$f'_\tau \partial_\tau \phi_{,\tau} = f''_\tau \phi_{,\tau}, \quad (60)$$

$$4\epsilon\phi_{,\tau} = f'_\tau \phi_{,4}. \quad (61)$$

Using the condition $\phi_{,4\tau} = 0$ and relation (61), one gets $\partial_{4\tau}\phi = 0$. If not all $f'_\tau \neq 0$, then, from obtained formulas it follows that

$$\phi = \frac{1}{2}a_1 \sum_{i=1}^6 f_i = a_1\varphi, \quad f_1 = f_2 = f_3 = f_4 = \epsilon x^4, \quad (62)$$

where φ is determined by formula (11). In the opposite case, when all $f'_\tau = 0$, we have

$$\begin{aligned} \phi_{,\tau} &= \phi_{,14} = \phi_{,23} = \phi_{,24} = \phi_{,\tau\tau} = 0, \\ \phi_{,34} &= -\frac{2\epsilon^2}{3A(f_\tau - f_4)}\left(\frac{b_{\tau\tau}}{g_{\tau\tau}} - \frac{b_{14}}{g_{14}}\right), \\ \phi_{,44} &= \partial_{44}\phi - \frac{2\epsilon(3\theta' - 2\epsilon^2 x^2)}{3A(f_\tau - f_4)}\left(\frac{b_{\tau\tau}}{g_{\tau\tau}} - \frac{b_{14}}{g_{14}}\right), \end{aligned} \quad (63)$$

where $\epsilon = 1$, since when $\epsilon = f'_\tau = 0$ the space is of constant curvature.

Put $(ijkl) = (4413), (4\tau 4\tau)$ in equation (20). One finds

$$b_{14}R_{413}^1 + b_{24}R_{413}^2 = g_{14}\phi_{,34}, \quad (64)$$

$$b_{14}R_{\tau 4\tau}^1 + b_{24}R_{\tau 4\tau}^2 + b_{\tau\tau}R_{44\tau}^\tau = -g_{\tau\tau}\phi_{,44}. \quad (65)$$

Expressing b_{24} from (64) and substituting the result in (65), taking into account (63), one obtains $\partial_{44}\phi = 0$ coming to the result needed.

Using equations (19) with $(ijk) = (166), (266), (366), (466), (165), (566), (666)$ in the case of the h -space of the [51] type, one gets

$$\begin{aligned} \phi_{,1} &= \phi_{,2} = \phi_{,3} = \phi_{,4} = 0, \\ \phi_{,6} &= \frac{1}{2}\frac{f'_6}{f_6 - f_5}\left(\frac{b_{66}}{g_{66}} - \frac{b_{15}}{g_{15}}\right), \quad \phi_{,5} = \frac{5}{2}\frac{\epsilon}{f_6 - f_5}\left(\frac{b_{66}}{g_{66}} - \frac{b_{15}}{g_{15}}\right), \\ \frac{\partial_6 b_{66}}{g_{66}} &= -5f'_6(f_5 - f_6)^{-1}\frac{b_{66}}{g_{66}} + 4\phi_{,6}, \end{aligned} \quad (66)$$

and, hence,

$$f'_6\partial_6\phi_{,6} = f''_6\phi_{,6}, \quad 5\epsilon\phi_{,6} = f'_6\phi_{,5}. \quad (67)$$

The last equation and the condition $\phi_{,56} = 0$ lead to

$$\partial_{56}\phi = 0. \quad (68)$$

Integrating equations (67) and (68) when $f'_6 \neq 0$, one obtains

$$\phi = \frac{1}{2}a_1 \sum_{i=1}^6 f_i = a_1\varphi, \quad f_1 = f_2 = f_3 = f_4 = f_5 = \epsilon x^5, \quad (69)$$

where φ is obtained by formula (11).

In the case of $f'_6 = 0$, from the formulas obtained we can find

$$\begin{aligned}\phi_{,6} &= \phi_{,15} = \phi_{,24} = \phi_{,25} = \phi_{,33} = \phi_{,34} = \phi_{,35} = \phi_{,66} = 0, \\ \phi_{,45} &= -\frac{15\epsilon^2}{16A(f_6 - f_5)}\left(\frac{b_{66}}{g_{66}} - \frac{b_{15}}{g_{15}}\right), \\ \phi_{,55} &= \partial_{55}\phi - \frac{5\epsilon(4\theta' - 3\epsilon^2x^3)}{4A(f_6 - f_5)}\left(\frac{b_{66}}{g_{66}} - \frac{b_{15}}{g_{15}}\right),\end{aligned}\tag{70}$$

where $\epsilon = 1$, since we consider the space of nonconstant curvature.

From (20) with $(ijkl) = (5514), (5656)$ and taking into account (19), one obtains the equality $\partial_{55}\phi = 0$, which leads to relation (69).

Therefore, the theorem is proved.

I am grateful to professor A.V.Aminova for constant attention to this work and for useful discussions.

References

- [A95] Aminova A. V. Lie algebras of infinitesimal projective transformations of Lorentz manifolds (in Russian) // Uspekhi Mat. Nauk **50** (1995), no. 1(301), 69-142.
- [E] L.P. Eisenhart, Riemannian geometry, Princeton University Press, 1997.
- [Z] Z. Zakirova, PhD thesis, 2001, Kazan State University
- [Z1] Z. Zakirova, First integral of the geodesics equations in 6d h-spaces // in: Proc. of Geometry Seminar, Kazan, 1996
- [Z2] Z. Zakirova, 6-dimensional h -spaces of the special type // Lobachevskiy Int. geom. seminar, Kazan, 1997: Abstracts of contributed papers – Kazan, 1997.- P. 52.
- [Z3] Z. Zakirova, On one solution of the Eisenhart equation. // "Latest problems of the field theory. 1999-2000". Editor A. V. Aminova.- Kazan, 2000. . 104-109.
- [Z4] Z. Zakirova, First integral of the geodesics equations in h -space of the [411] type. // Izv. Vuzov. Mathematics. N9 (448), 1999, . 78-79.
- [Z5] Z.Zakirova, On projective group properties of the 6d h -space of the type [33], math.DG/0101052; submitted to "Classical & Quantum Gravity"
- [Z6] Z.Zakirova, Rigid 6-dimensional h -spaces of constant curvature, math.DG/0301105